

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 2 of 33 of February 12, 2010 Amendment


Amendments to the Claims

This listing of claims will replace all prior versions, and listings of claims in the application.

Listing of the Claims

Claims 1-48 (Cancelled)

49. (Withdrawn) A method for the treatment of inflammation which comprises administering to an individual in need thereof, an effective amount of a compound of the general formula I:



or of a pharmaceutically acceptable salt thereof, wherein R1 is C₁₂-C₂₄ alkyl or C₁₀-C₂₄ alkenyl, and A is a residue containing at least one acyclic or cyclic amino group and/or at least one heteroaromatic ring containing a tertiary or quaternary nitrogen atom.

50. (Withdrawn) The method according to claim 49 wherein R1 is a C₁₂-C₂₀ alkyl or alkenyl.

51. (Withdrawn) The method according to claim 50 wherein R1 is a C₁₆-C₁₈ alkyl or alkenyl.

52. (Withdrawn) The method according to claim 51 wherein R1 is hexadecyl, octadecyl, hexadecenyl, octadecenyl, cis-9-octadecenyl, trans-9-octadecenyl, cis-9,12-octadecadienyl,


Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 3 of 33 of February 12, 2010 Amendment

cis-6,9,12-octadecatrienyl, or cis-9,12,15-octadecatrienyl;

53. (Withdrawn) The method according to claim 52 wherein R1 is cis-9-octadecenyl or trans-9-octadecenyl.

54. (Withdrawn) The method according to claim 49 wherein in said compound of formula I the residue A is selected from the group consisting of:


(i)



wherein R2 is H, C₁-C₆ alkyl, aryl or aralkyl, wherein any aryl moiety may be unsubstituted or substituted by nitro, cyano, halo, hydroxy, NR₆R₇, or CR₈R₈NR₆R₇; R3 is H, a pair of electrons or C₁-C₆ alkyl; R4 and R5 each independently is H or C₁-C₆ alkyl, or R4 and R5 together with the nitrogen atom to which they are attached form a 5-7 membered saturated ring optionally interrupted by an oxygen atom or by a nitrogen atom optionally substituted by C₁-C₆ alkyl; and R6, R7 and R8 each independently is H or C₁-C₆ alkyl;

(ii) phenyl substituted by NR₆R₇ or CR₈R₈NR₆R₇, wherein R6, R7 and R8 each is independently H or C₁-C₆ alkyl; and


(iii)



Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 4 of 33 of February 12, 2010 Amendment

wherein R9 is H, C₁-C₆ alkyl or indolyl(C₁-C₄)alkyl, and X⁻ is a counter ion, or R9 is a pair of electrons and X is absent.

55. (Withdrawn) The method according to claim 54 wherein the residue A is of the formula:



wherein R₂ is H; a straight or branched C₁-C₆ alkyl selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, pentyl and hexyl; phenyl, benzyl or p-hydroxybenzyl; R₃ is H, a pair of electrons or C₁-C₆ alkyl; R₄ and R₅ is each independently H or C₁-C₆ alkyl, or R₄ and R₅ together with the nitrogen atom to which they are attached form a 5-7 membered saturated ring optionally interrupted by an oxygen atom or by a nitrogen atom optionally substituted by C₁-C₆ alkyl, said ring being selected from the group consisting of pyrrolidine, piperidine, morpholine, piperazine, and 4-methylpiperazine.

56. (Withdrawn) The method according to claim 55 wherein R₂ is H or phenyl, R₃ is H or a pair of electrons, R₄ and R₅ are each H or C₁-C₆ alkyl, or R₄ and R₅ together with the N atom to which they are attached form a morpholine ring or a piperazine ring optionally substituted at the nitrogen atom at position 1 or 4 by methyl.

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 5 of 33 of February 12, 2010 Amendment

57. (Withdrawn) The method according to claim 56 wherein said compound is selected from the group consisting of:

N,N-Dimethylamino-acetic acid octadec-(Z)-9-enyl ester;
(4-Methyl-piperazin-1-yl)-acetic acid octadec-(Z)-9-enyl ester tartrate;
(4-Methyl-piperazin-1-yl)-acetic acid octadecyl ester tartrate;
4-Methyl-4-octadec-(Z)-9-enyloxycarbonylmethyl-morpholin-4-ium chloride;
 α -Amino- α -phenyl-acetic acid octadec-(Z)-9-enyl ester HCl salt; and
Piperazin-1-yl-acetic acid octadec-(Z)-9-enyl ester bitartrate.


58. (Withdrawn) The method according to claim 54 wherein the residue A is phenyl substituted by NR₆R₇ or CR₈R₈NR₆R₇, wherein R₆, R₇ and R₈ is each independently H or C₁-C₆ alkyl.

59. (Withdrawn) The method according to claim 58 wherein A is phenyl substituted by CR₈R₈NR₆R₇, wherein R₈ is H and R₆ and R₇ is each methyl.

60. (Withdrawn) The method according to claim 59, wherein said compound is 4-dimethylaminomethyl-benzoic acid octadec-(Z)-9-enyl ester HCl or 4-dimethylaminomethyl-benzoic acid octadec-(E)-9-enyl ester HCl.

61. (Withdrawn) The method according to claim 54 wherein A is the group:

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 6 of 33 of February 12, 2010 Amendment



wherein R9 is H, C₁-C₆ alkyl or indolyl(C₁-C₄)alkyl and X⁻ is a counter ion, or R9 is a pair of electrons and X⁻ is absent.

62. (Withdrawn) The method according to claim 61 wherein R9 is a pair of electrons and X⁻ is absent, or R9 is methyl or indolyethyl and X⁻ is a counter ion selected from the group consisting of chloride, bromide, iodide and tosylate.

63. (Withdrawn) The method according to claim 62 wherein said compound is selected from the group consisting of:

Nicotinic acid octadec-(Z)-9-enyl ester;
1-Methyl-3-octadec-(Z)-9-enyloxycarbonyl-pyridinium iodide;
1-Methyl-3-octadec-(Z)-9-enyloxycarbonyl-pyridinium chloride;
1-Methyl-3-octadec-(Z)-9-enyloxycarbonyl-pyridinium tosylate; and
1-[(2-(1H-indol-3-yl)-ethyl)-3-octadec-(Z)-9-enyloxycarbonyl-pyridinium bromide.

64. (Withdrawn) The method according to claim 49 for treatment of immunologically-mediated inflammation.

65. (Withdrawn) The method according to claim 64 for the treatment of an immunologically-mediated chronic or acute inflammatory disease, disorder or condition.

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 7 of 33 of February 12, 2010 Amendment

66. (Withdrawn) The method according to claim 65 for the treatment of an autoimmune disease, a severe allergy, asthma, or an inflammation associated with a disease, disorder or condition selected from graft rejection, a chronic degenerative disease such as Alzheimer's disease, neuroprotection, organ regeneration, chronic ulcers of the skin, or schizophrenia.

67. (Withdrawn) The method according to claim 66 wherein said autoimmune disease, disorder or condition is multiple sclerosis or a human arthritic condition.

68. (Withdrawn) The method according to claim 67 wherein said human arthritic condition is rheumatoid arthritis, reactive arthritis with Reiter's syndrome, ankylosing spondylitis or other inflammation of the joints mediated by the immune system.

69. (Withdrawn) The method according to claim 65 wherein said immunologically-mediated inflammatory disease, disorder or condition is myasthenia gravis, Guillain-Barré syndrome, or other inflammatory disease of the nervous system; psoriasis, pemphigus vulgaris or other diseases of the skin; systemic lupus erythematosus, glomerulonephritis or other disease affecting the kidneys; atherosclerosis or other inflammation of the blood vessels; autoimmune hepatitis, inflammatory bowel diseases, pancreatitis, or other disorder of the gastrointestinal system; type 1 diabetes mellitus, autoimmune thyroiditis, or other disease of the endocrine system.

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 8 of 33 of February 12, 2010 Amendment


70. (Withdrawn) The method according to claim 69 wherein said immunologically-mediated inflammatory disease or disorder is psoriasis.

71. (Withdrawn) The method according to claim 49 wherein said compound is administered by oral, topical, intradermal or parenteral route.

72. (Withdrawn) The method according to claim 71 wherein said compound is administered by subcutaneous, intravenous, or intramuscular route.

Claims 73-134. (Cancelled)

135. (Withdrawn) A method of treating a T-cell mediated disease, disorder or condition, which comprises administering to an individual in need an effective amount of a therapeutic preparation comprising an antigen recognized by inflammatory T cells associated with the pathogenesis of said T-cell mediated disease, disorder or condition, and an adjuvant of the general formula Ia:



[Ia]

or a pharmaceutically acceptable salt thereof, wherein R1 is C10-C24 alkyl or C10-C24 alkenyl, and A is a residue containing at least one acyclic or cyclic amino group and/or at least one heteroaromatic ring containing a tertiary or quaternary nitrogen atom, but excluding the compounds wherein

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 9 of 33 of February 12, 2010 Amendment

R1 is C18 alkyl and A is a residue containing at least one acyclic amino group or -CO-A is the residue of proline.

136. (Withdrawn) A method of causing a shifting of T-cell cytokine response from T_H1 to T_H2 in an individual suffering from a T-cell mediated disease, disorder or condition, which comprises administering to said individual in need an effective amount of a therapeutic preparation comprising an antigen recognized by inflammatory T cells associated with the pathogenesis of said T-cell mediated disease, disorder or condition and an adjuvant of the general formula Ia:



or a pharmaceutically acceptable salt thereof, wherein R1 is C10-C24 alkyl or C10-C24 alkenyl, and A is a residue containing at least one acyclic or cyclic amino group and/or at least one heteroaromatic ring containing a tertiary or quaternary nitrogen atom, but excluding the compounds wherein R1 is C18 alkyl and A is a residue containing at least one acyclic amino group or -CO-A is the residue of proline.

137. (Withdrawn) The method according to claim 136 wherein said therapeutic preparation causes a decrease in IL-2 or IFN- γ T-cell cytokine response and an increase in IL-4 or IL-10 T-cell cytokine response.

138. (Withdrawn) The method according to claim 135 wherein R1

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 10 of 33 of February 12, 2010 Amendment

is a C₁₂-C₂₀ alkyl or alkenyl.


139. (Withdrawn) The method according to claim 138 wherein R1 is a C₁₆-C₁₈ alkyl or alkenyl.

140. (Withdrawn) The method according to claim 139 wherein R1 is hexadecyl, octadecyl, hexadecenyl, octadecenyl, cis-9-octadecenyl, trans-9-octadecenyl, cis-9,12-octadecadienyl, cis-6,9,12-octadecatrienyl, or cis-9,12,15-octadecatrienyl.

141. (Withdrawn) The method according to claim 140 wherein R1 is cis-9-octadecenyl or trans-9-octadecenyl.

142. (Withdrawn) The method according to claim 135 wherein in said compound of formula Ia the residue A is selected from the group consisting of:

(i)




wherein R2 is H, C₁-C₆ alkyl, aryl or aralkyl, wherein any aryl moiety may be unsubstituted or substituted by nitro, cyano, halo, hydroxy, NR₆R₇, or CR₈R₈NR₆R₇; R3 is H, a pair of electrons or C₁-C₆ alkyl; R4 and R5 is each independently H or C₁-C₆ alkyl, or R4 and R5 together with the nitrogen atom to which they are attached form a 5-7 membered saturated ring optionally interrupted by an oxygen atom or by a nitrogen atom optionally substituted by C₁-C₆ alkyl, provided that R4 and R5

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 11 of 33 of February 12, 2010 Amendment

are not H or C₁-C₆ alkyl when R₁ is octadecyl; and R₆, R₇ and R₈ is each independently H or C₁-C₆ alkyl;


(ii) phenyl substituted by NR₆R₇ or CR₈R₈NR₆R₇, wherein R₆, R₇ and R₈ each is independently H or C₁-C₆ alkyl; and

(iii)



wherein R₉ is H, C₁-C₆ alkyl or indolyl(C₁-C₄)alkyl, and X⁻ is a counter ion, or R₉ is a pair of electrons and X is absent.

143. (Withdrawn) The method according to claim 142 wherein the residue A has the formula:



wherein R₂ is H; a straight or branched C₁-C₆ alkyl selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, pentyl and hexyl; phenyl, benzyl or p-hydroxybenzyl; R₃ is H, a pair of electrons or C₁-C₆ alkyl; R₄ and R₅ is each independently H or C₁-C₆ alkyl, or R₄ and R₅ together with the nitrogen atom to which they are attached form a 5-7 membered saturated ring optionally interrupted by an oxygen atom or by a nitrogen atom

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 12 of 33 of February 12, 2010 Amendment

optionally substituted by C₁-C₆ alkyl, said ring being selected from the group consisting of pyrrolidine, piperidine, morpholine, piperazine, and 4-methylpiperazine, provided that R₄ and R₅ are not H or C₁-C₆ alkyl when R₁ is octadecyl.

144. (Withdrawn) The method according to claim 143 wherein R₂ is H or phenyl, R₃ is H or a pair of electrons, R₄ and R₅ is each H or C₁-C₆ alkyl, or R₄ and R₅ together with the N atom to which they are attached form a morpholine ring or a piperazine ring optionally substituted at the nitrogen atom at position 1 or 4 by methyl, provided that R₄ and R₅ are not H or C₁-C₆ alkyl when R₁ is octadecyl.

145. (Withdrawn) The method according to claim 144 wherein said adjuvant is selected from the group consisting of:


N,N-Dimethylamino-acetic acid octadec-(Z)-9-enyl ester;
(4-Methyl-piperazin-1-yl)-acetic acid octadec-(Z)-9-enyl ester tartrate;
4-Methyl-4-octadec-(Z)-9-enyloxycarbonylmethyl-morpholin-4-ium chloride;
 α -Amino- α -phenyl-acetic acid octadec-(Z)-9-enyl ester HCl salt; and
Piperazin-1-yl-acetic acid octadec-(Z)-9-enyl ester bitartrate.

146. (Withdrawn) The method according to claim 135 wherein R₁ is C₁₈ alkyl and A is a residue containing at least one cyclic amino group and/or at least one heteroaromatic ring containing a tertiary or quaternary nitrogen atom, but excluding the compound wherein -CO-A is the residue of proline.

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 13 of 33 of February 12, 2010 Amendment

147. (Withdrawn) The method according to claim 146 wherein the residue A has the formula:

(i)



wherein R2 is H, C₁-C₆ alkyl, aryl or aralkyl, wherein any aryl moiety may be unsubstituted or substituted by nitro, cyano, halo, hydroxy, NR₆R₇, or CR₈R₈NR₆R₇; R3 is H, a pair of electrons or C₁-C₆ alkyl; R4 and R5 together with the nitrogen atom to which they are attached form a 5-7 membered saturated ring optionally interrupted by an oxygen atom or by a nitrogen atom optionally substituted by C₁-C₆ alkyl; and R6, R7 and R8 is each independently H or C₁-C₆ alkyl.

148. (Withdrawn) The method according to claim 147 wherein R2 is H or phenyl, R3 is H or a pair of electrons, and R4 and R5 together with the N atom to which they are attached form a morpholine ring or a piperazine ring optionally substituted at the nitrogen atom at position 4 by methyl.

149. (Withdrawn) The method according to claim 148 wherein said adjuvant is (4-methyl-piperazin-1-yl)-acetic acid octadecyl ester tartrate.


150. (Withdrawn) The method according to claim 142 wherein A is phenyl substituted by NR₆R₇ or CR₈R₈NR₆R₇, wherein R6, R7 and R8 is each independently H or 1 C₁-C₆ alkyl.

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 14 of 33 of February 12, 2010 Amendment

151. (Withdrawn) The method according to claim 150 wherein A is phenyl substituted by CR₈R₈NR₆R₇, wherein R₈ is H and R₆ and R₇ is each methyl.

152. (Withdrawn) The method according to claim 151 wherein said adjuvant is 4-dimethylaminomethyl-benzoic acid octadec-(Z)-9-enyl ester HCl or 4-dimethylaminomethyl-benzoic acid octadec-(E)-9-enyl ester HCl.

153. (Withdrawn) The method according to claim 142 wherein A is the group:



wherein R₉ is H, C₁-C₆ alkyl or indolyl(C₁-C₄)alkyl and X⁻ is a counter ion, or R₉ is a pair of electrons and X⁻ is absent.

154. (Withdrawn) The method according to claim 153 wherein R₉ is a pair of electrons and X⁻ is absent, or R₉ is methyl or indolyethyl and X⁻ is a counter ion selected from the group consisting of chloride, bromide, iodide and tosylate.

155. (Withdrawn) The method according to claim 154 wherein said adjuvant is selected from the group consisting of:

Nicotinic acid octadec-(Z)-9-enyl ester;

1-Methyl-3-octadec-(Z)-9-enyloxycarbonyl-pyridinium iodide;

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 15 of 33 of February 12, 2010 Amendment

1-Methyl-3-octadec-(Z)-9-enyloxycarbonyl-pyridinium
chloride;
1-Methyl-3-octadec-(Z)-9-enyloxycarbonyl-pyridinium
tosylate; and
1-[(2-(1H-indol-3-yl)-ethyl)-3-octadec-(Z)-9-
enyloxycarbonyl-pyridinium bromide.

156. (Withdrawn) The method according to claim 135 wherein
said antigen raises a humoral response in said individual.

157. (Withdrawn) The method according to claim 135 wherein
said antigen raises a cellular response in said individual.

158. (Withdrawn) The method according to claim 135 wherein said
T-cell mediated disease is an autoimmune disease and said
antigen is a peptide.

159. (Withdrawn) The method according to claim 158, wherein
said autoimmune disease is an organ-specific autoimmune
disease.

160. (Withdrawn) The method according to claim 159 wherein said
organ-specific autoimmune disease is type I diabetes mellitus,
multiple sclerosis, rheumatoid arthritis or autoimmune
thyroiditis.

161. (Withdrawn) The method according to claim 160 for the
treatment of multiple sclerosis wherein said antigen is a
peptide derived from the sequence of myelin basic protein
(MBP) or an analogue thereof that is recognized by T-cells
involved in the pathogenesis of multiple sclerosis.


Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 16 of 33 of February 12, 2010 Amendment

162. (Withdrawn) The method according to claim 160 for the treatment of multiple sclerosis wherein said antigen is a copolymer recognized by T-cells involved in the pathogenesis of multiple sclerosis.

163. (Withdrawn) The method according to claim 162 wherein said antigen is glatiramer acetate.


164. (Withdrawn) The method according to claim 135 wherein said therapeutic preparation comprises said adjuvant and an antigen useful for treatment of an autoimmune disease, a neurodegenerative disease such as Alzheimer's disease or Parkinson disease, a cancer such as melanoma, or an infectious disease such as a bacterial or viral infection.

165. (Currently amended) A compound of the general formula:



wherein

(i) R1 is C₂₀-C₂₄ alkyl or C₁₀-C₂₄ alkenyl and A is a residue of the formula:




wherein R2 is H, C₁-C₆ alkyl, aryl, or aralkyl, wherein any aryl moiety may be unsubstituted or substituted by nitro, cyano, halo, hydroxy, NR₆R₇, or CR₈R₈NR₆R₇; R3 is H, a pair of electrons, or C₁-C₆ alkyl; R4 and R5 each independently is H or C₁-C₆ alkyl, or R4 and R5 together with the nitrogen atom to

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 17 of 33 of February 12, 2010 Amendment

which they are attached form a 5-7 membered saturated ring optionally interrupted by an oxygen atom or by a nitrogen atom optionally substituted by C₁-C₆ alkyl; and R₆, R₇ and R₈ each independently is H or C₁-C₆ alkyl,


wherein when R₃ is a pair of electrons and R₄ and R₅ are each H, R₂ is C₅-C₆ alkyl, aryl, or aralkyl, wherein any aryl moiety may be unsubstituted or substituted by nitro, cyano, halo, hydroxy, NR₆R₇, or CR₈R₈NR₆R₇; or

(ii) R₁ is C₁₈ alkyl and A is a residue of the formula:



wherein R₂ is H; R₃ is a pair of electrons; and R₄ and R₅ together with the nitrogen atom to which they are attached form a 5-7 membered saturated ring optionally interrupted by an oxygen atom or by a nitrogen atom optionally substituted by C₁-C₆ alkyl; or

(iii) R₁ is C₁₂-C₁₆ alkyl and A is a residue of the formula:




wherein R₂ is unsubstituted aryl, or aryl or aralkyl wherein the aryl moiety is substituted by nitro, cyano, halo, hydroxy, NR₆R₇, or CR₈R₈NR₆R₇; R₃ is H, a pair of electrons, or C₁-C₆ alkyl; R₄ and R₅ each independently is H or C₁-C₆ alkyl, or R₄ and R₅ together with the nitrogen atom to which they are attached form a 5-7 membered saturated ring optionally interrupted by an oxygen atom or by a nitrogen atom

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 18 of 33 of February 12, 2010 Amendment

optionally substituted by C₁-C₆ alkyl; and R₆, R₇ and R₈ each independently is H or C₁-C₆ alkyl; or

(iv) R₁ is C₁₀ alkyl and A is a residue of the formula:




wherein R₂ is C₁-C₆ alkyl; R₃ is H, a pair of electrons, or C₁-C₆ alkyl; R₄ and R₅ each independently is H or C₁-C₆ alkyl, or R₄ and R₅ together with the nitrogen atom to which they are attached form a 5-7 membered saturated ring optionally interrupted by an oxygen atom or by a nitrogen atom optionally substituted by C₁-C₆ alkyl; and R₆, R₇ and R₈ each independently is H or C₁-C₆ alkyl,

wherein when R₃ is a pair of electrons and R₄ and R₅ are each H, R₂ is C₅-C₆ alkyl, aryl, or aralkyl, wherein any aryl moiety may be unsubstituted or substituted by nitro, cyano, halo, hydroxy, NR₆R₇, or CR₈R₈NR₆R₇; or

(v) R₁ is C₁₀-C₂₄ alkyl or C₁₀-C₂₄ alkenyl and A is phenyl substituted by NR₆R₇ or CR₈R₈NR₆R₇, wherein R₆, R₇ and R₈ each independently is H or C₁-C₆ alkyl, but excluding the compounds wherein R₁ is C₁₀-C₁₆ alkyl and A is phenyl substituted by -CH₂-NH₂; or

(vi) R₁ is C₁₀-C₂₄ alkyl or C₁₀-C₂₄ alkenyl and A is a group of the formula:



wherein R₉ is C₁-C₆ alkyl or indolyl(C₁-C₆)alkyl and X⁻ is a

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 19 of 33 of February 12, 2010 Amendment

counter ion; and pharmaceutically acceptable salts thereof.

166. (Original) A compound according to claim 165(i), (v) or (vi) wherein R1 is a C₁₂-C₁₈ alkenyl.

167. (Original) A compound according to claim 166 wherein R1 is a C₁₆-C₁₈ alkenyl.

168. (Original) A compound according to claim 167 wherein R1 is cis-9-octadecenyl or trans-9-octadecenyl.

169. (Withdrawn) A compound according to claim 168 wherein R2 is H or phenyl, R3 is H or a pair of electrons, and R4 and R5 are methyl or together with the N atom to which they are attached form a morpholino or a piperazine ring optionally substituted at the nitrogen atom at position 4 by methyl.

170. (Withdrawn) A compound according to claim 169 selected from the group consisting of:

N,N-Dimethylamino-acetic acid octadec-(Z)-9-enyl ester;

(4-Methyl-piperazin-1-yl)-acetic acid octadec-(Z)-9-enyl ester tartrate;

4-Methyl-4-octadec-(Z)-9-enyloxycarbonylmethyl-morpholin-4-ium chloride;

Piperazin-1-yl-acetic acid octadec-(Z)-9-enyl ester bitartrate.

171. (Withdrawn) A compound according to claim 165(ii) wherein R1 is octadecyl and R4 and R5 together with the N atom to which they are attached form a morpholino or a piperazine ring optionally substituted at the nitrogen atom at position 4 by methyl.

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 20 of 33 of February 12, 2010 Amendment

172. (Withdrawn) A compound according to claim 171 which is (4-methyl-piperazin-1-yl)-acetic acid octadecyl ester tartrate.

173. (Withdrawn) A compound according to claim 165(vi) wherein R1 is C₁₂-C₂₀ alkyl or C₁₂-C₂₀ alkenyl and A is phenyl substituted by CR₈R₈NR₆R₇, wherein R₈ is H and R₆ and R₇ is each H or C₁-C₆ alkyl.

174. (Withdrawn) A compound according to claim 173 wherein R1 is C₁₆-C₁₈ alkenyl.

175. (Withdrawn) A compound according to claim 174 wherein R1 is cis-9-octadecenyl or trans-9-octadecenyl.

176. (Withdrawn) A compound according to claim 175 which is 4-dimethylaminomethyl-benzoic acid octadec-(Z)-9-enyl ester HCl or 4-dimethylaminomethyl-benzoic acid octadec-(E)-9-enyl ester HCl.

177. (Withdrawn) A compound according to claim 165(vi) wherein R1 is C₁₂-C₂₀ alkyl or C₁₂-C₂₀ alkenyl.

178. (Withdrawn) A compound according to claim 177 wherein R1 is C₁₆-C₁₈ alkyl or alkenyl.

179. (Withdrawn) A compound according to claim 178 wherein R1 is cis-9-octadecenyl or trans-9-octadecenyl.

180. (Withdrawn) A compound according to claim 179 selected

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 21 of 33 of February 12, 2010 Amendment

from the group consisting of:

1-Methyl-3-octadec-(Z)-9-enyloxycarbonyl-pyridinium
iodide;
1-Methyl-3-octadec-(Z)-9-enyloxycarbonyl-pyridinium
chloride;
1-Methyl-3-octadec-(Z)-9-enyloxycarbonyl-pyridinium
tosylate; and
1-[(2-(1H-indol-3-yl)-ethyl)-3-octadec-(Z)-9-
enyloxycarbonyl-pyridinium bromide.

181. (Currently amended) A compound according to claim 165(i) wherein R1 is cis-9-octadecenyl, R2 is phenyl, R3 is a pair of electrons and R4 and R5 is each H.

182. (Original) The compound of claim 181 which is α -amino- α -phenyl-acetic acid octadec-(Z)-9-enyl ester HCl salt.

183. (Previously presented) A pharmaceutical composition comprising a compound according to claim 165 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

184. (Previously presented) A pharmaceutical composition according to claim 183 for the treatment of an-immunologically mediated acute or chronic inflammatory disease, disorder or condition.

185. (Previously presented) A therapeutic composition comprising an antigen and an adjuvant according to claim 165.

186. (Withdrawn) The method according to claim 136 wherein R1

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 22 of 33 of February 12, 2010 Amendment

is a C₁₂-C₂₀ alkyl or alkenyl.


187. (Withdrawn) The method according to claim 186 wherein R1 is a C₁₆-C₁₈ alkyl or alkenyl.

188. (Withdrawn) The method according to claim 187 wherein R1 is hexadecyl, octadecyl, hexadecenyl, octadecenyl, cis-9-octadecenyl, trans-9-octadecenyl, cis-9,12-octadecadienyl, cis-6,9,12-octadecatrienyl, or cis-9,12,15-octadecatrienyl,

189. (Withdrawn) The method according to claim 188 wherein R1 is cis-9-octadecenyl or trans-9-octadecenyl.

190. (Withdrawn) The method according to claim 136 wherein in said compound of formula Ia the residue A is selected from the group consisting of:

(i)




wherein R2 is H, C₁-C₆ alkyl, aryl or aralkyl, wherein any aryl moiety may be unsubstituted or substituted by nitro, cyano, halo, hydroxy, NR₆R₇, or CR₈R₈NR₆R₇; R3 is H, a pair of electrons or C₁-C₆ alkyl; R4 and R5 is each independently H or C₁-C₆ alkyl, or R4 and R5 together with the nitrogen atom to which they are attached form a 5-7 membered saturated ring optionally interrupted by an oxygen atom or by a nitrogen atom optionally substituted by C₁-C₆ alkyl, provided that R4 and R5

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 23 of 33 of February 12, 2010 Amendment

are not H or C₁-C₆ alkyl when R₁ is octadecyl; and R₆, R₇ and R₈ is each independently H or C₁-C₆ alkyl;


(ii) phenyl substituted by NR₆R₇ or CR₈R₈NR₆R₇, wherein R₆, R₇ and R₈ each is independently H or C₁-C₆ alkyl; and

(iii)



wherein R₉ is H, C₁-C₆ alkyl or indolyl(C₁-C₄)alkyl, and X⁻ is a counter ion, or R₉ is a pair of electrons and X is absent.

191. (Withdrawn) The method according to claim 190 wherein the residue A has the formula:



wherein R₂ is H; a straight or branched C₁-C₆ alkyl selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, pentyl and hexyl; phenyl, benzyl or p-hydroxybenzyl; R₃ is H, a pair of electrons or C₁-C₆ alkyl; R₄ and R₅ is each independently H or C₁-C₆ alkyl, or R₄ and R₅ together with the nitrogen atom to which they are attached form a 5-7 membered saturated ring optionally interrupted by an oxygen atom or by a nitrogen atom

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 24 of 33 of February 12, 2010 Amendment

optionally substituted by C₁-C₆ alkyl, said ring being selected from the group consisting of pyrrolidine, piperidine, morpholine, piperazine, and 4-methylpiperazine, provided that R₄ and R₅ are not H or C₁-C₆ alkyl when R₁ is octadecyl.

192. (Withdrawn) The method according to claim 191 wherein R₂ is H or phenyl, R₃ is H or a pair of electrons, R₄ and R₅ is each H or C₁-C₆ alkyl, or R₄ and R₅ together with the N atom to which they are attached form a morpholine ring or a piperazine ring optionally substituted at the nitrogen atom at position 1 or 4 by methyl, provided that R₄ and R₅ are not H or C₁-C₆ alkyl when R₁ is octadecyl.

193. (Withdrawn) The method according to claim 192 wherein said adjuvant is selected from the group consisting of:


N,N-Dimethylamino-acetic acid octadec-(Z)-9-enyl ester;
(4-Methyl-piperazin-1-yl)-acetic acid octadec-(Z)-9-enyl ester tartrate;
4-Methyl-4-octadec-(Z)-9-enyloxycarbonylmethyl-morpholin-4-ium chloride;
 α -Amino- α -phenyl-acetic acid octadec-(Z)-9-enyl ester HCl salt; and
Piperazin-1-yl-acetic acid octadec-(Z)-9-enyl ester bitartrate.

194. (Withdrawn) The method according to claim 136 wherein R₁ is C₁₈ alkyl and A is a residue containing at least one cyclic amino group and/or at least one heteroaromatic ring containing a tertiary or quaternary nitrogen atom, but excluding the compound wherein -CO-A is the residue of proline.

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 25 of 33 of February 12, 2010 Amendment

195. (Withdrawn) The method according to claim 194 wherein the residue A has the formula:

(i)



wherein R2 is H, C₁-C₆ alkyl, aryl or aralkyl, wherein any aryl moiety may be unsubstituted or substituted by nitro, cyano, halo, hydroxy, NR₆R₇, or CR₈R₈NR₆R₇; R3 is H, a pair of electrons or C₁-C₆ alkyl; R4 and R5 together with the nitrogen atom to which they are attached form a 5-7 membered saturated ring optionally interrupted by an oxygen atom or by a nitrogen atom optionally substituted by C₁-C₆ alkyl; and R6, R7 and R8 is each independently H or C₁-C₆ alkyl.

196. (Withdrawn) The method according to claim 195 wherein R2 is H or phenyl, R3 is H or a pair of electrons, and R4 and R5 together with the N atom to which they are attached form a morpholine ring or a piperazine ring optionally substituted at the nitrogen atom at position 4 by methyl.

197. (Withdrawn) The method according to claim 196 wherein said adjuvant is (4-methyl-piperazin-1-yl)-acetic acid octadecyl ester tartrate.


198. (Withdrawn) The method according to claim 197 wherein A is phenyl substituted by NR₆R₇ or CR₈R₈NR₆R₇, wherein R6, R7 and R8 is each independently H or 1 C₁-C₆ alkyl.

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 26 of 33 of February 12, 2010 Amendment

199. (Withdrawn) The method according to claim 198 wherein A is phenyl substituted by CR₈R₈NR₆R₇, wherein R₈ is H and R₆ and R₇ is each methyl.

200. (Withdrawn) The method according to claim 199 wherein said adjuvant is 4-dimethylaminomethyl-benzoic acid octadec-(Z)-9-enyl ester HCl or 4-dimethylaminomethyl-benzoic acid octadec-(E)-9-enyl ester HCl.

201. (Withdrawn) The method according to claim 190 wherein A is the group:



wherein R₉ is H, C₁-C₆ alkyl or indolyl(C₁-C₄)alkyl and X⁻ is a counter ion, or R₉ is a pair of electrons and X⁻ is absent.

202. (Withdrawn) The method according to claim 201 wherein R₉ is a pair of electrons and X⁻ is absent, or R₉ is methyl or indolyethyl and X⁻ is a counter ion selected from the group consisting of chloride, bromide, iodide and tosylate.

203. (Withdrawn) The method according to claim 202 wherein said adjuvant is selected from the group consisting of:

Nicotinic acid octadec-(Z)-9-enyl ester;
1-Methyl-3-octadec-(Z)-9-enyloxycarbonyl-pyridinium
iodide;

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 27 of 33 of February 12, 2010 Amendment

1-Methyl-3-octadec-(Z)-9-enyloxycarbonyl-pyridinium
chloride;
1-Methyl-3-octadec-(Z)-9-enyloxycarbonyl-pyridinium
tosylate; and
1-[(2-(1H-indol-3-yl)-ethyl)-3-octadec-(Z)-9-
enyloxycarbonyl-pyridinium bromide.

204. (Withdrawn) The method according to claim 136 wherein
said antigen raises a humoral response in said individual.

205. (Withdrawn) The method according to claim 136 wherein
said antigen raises a cellular response in said individual.

206. (Withdrawn) The method according to claim 136 wherein said
T-cell mediated disease is an autoimmune disease and said
antigen is a peptide.

207. (Withdrawn) The method according to claim 206, wherein
said autoimmune disease is an organ-specific autoimmune
disease.

208. (Withdrawn) The method according to claim 207 wherein said
organ-specific autoimmune disease is type I diabetes mellitus,
multiple sclerosis, rheumatoid arthritis or autoimmune
thyroiditis.

209. (Withdrawn) The method according to claim 208 for the
treatment of multiple sclerosis wherein said antigen is a
peptide derived from the sequence of myelin basic protein
(MBP) or an analogue thereof that is recognized by T-cells
involved in the pathogenesis of multiple sclerosis.

Applicants: Meir Shinitzky et al.
Serial No.: 10/530,776
Filing Date: December 19, 2005
Page 28 of 33 of February 12, 2010 Amendment

210. (Withdrawn) The method according to claim 209 for the treatment of multiple sclerosis wherein said antigen is a copolymer recognized by T-cells involved in the pathogenesis of multiple sclerosis.

211. (Withdrawn) The method according to claim 210 wherein said antigen is glatiramer acetate.

212. (Withdrawn) The method according to claim 136 wherein said therapeutic preparation comprises said adjuvant and an antigen useful for treatment of an autoimmune disease, a neurodegenerative disease such as Alzheimer's disease or Parkinson disease, a cancer such as melanoma, or an infectious disease such as a bacterial or viral infection.